



LIBRARY OF THE
UNIVERSITY OF ILLINOIS
AT URBANA-CHAMPAIGN

590.5

FI

v. 39

cop. 3



NATURAL HISTORY,
SURVEY

• 1. 1. 1.

• 2. 2. 2.

• 3. 3. 3.

• 4. 4. 4.

9
12
3

FIELDIANA • ZOOLOGY

Published by
CHICAGO NATURAL HISTORY MUSEUM

Volume 39

FEBRUARY 28, 1958

No. 12

THE RACES OF THE BUSH SHRIKE

Dryoscopus cubla

AUSTIN L. RAND

CHIEF CURATOR, DEPARTMENT OF ZOOLOGY

The bush shrike, *Dryoscopus cubla*, is a bird of brush and woodland of South and eastern Africa. Chapin (1954, Bull. Amer. Mus. Nat. Hist., 75B: 63) and Mackworth-Praed and Grant (1955, Bds. East and N. E. Afr., 2: 617) have reviewed the species, with some conflicting views, and it appears that an additional subspecies must be recognized. It is possible that *cubla* may prove to be conspecific with *gambensis* of northern Africa, but I prefer to keep them separate.

The races I recognize are as follows:

***Dryoscopus cubla affinis* Gray, 1837**

The bird of coastal East Africa (Juba River–Dar-es-Salaam) with few or no white edgings to the wings.

Wing measurements.—Juba River, 76–81; coastal Kenya, 75–84; Zanzibar, 77 mm.

This is sometimes kept as a separate species but van Someren's discussion (1932, Nov. Zool., 37: 309) leaves little doubt that it is conspecific with *cubla* despite irregularities in intergradation, and our specimens support this view.

***Dryoscopus cubla nairobiensis*, new subspecies**

Type.—Chicago Natural History Museum no. 116606 from Nairobi, Kenya, 5400 feet altitude. Adult female, collected June 7, 1908; F. J. Jackson collection.

Diagnosis.—Like *D. c. affinis* in that the female possesses a black spot in front of the eye, but differs in the male, which has the rump more tinged with gray, and differs in both sexes in having more white in the edgings of the secondaries and in the upper wing coverts, and

Library of Congress Catalog Card Number: 58-9214

being slightly larger. Differs from all the other races in having a black spot in front of the eye (not with lores all white) in the female.

Wing measurements.—Kenya: Kijabe, ♂ 85, 85; Nairobi, ♂ (10) 81–86 (av. 83.4); ♀ (8) 77–81 (av. 79.2); Ukamba, ♂ 80, 82; ♀ 80. Tanganyika Territory: Kilimanjaro, ♂ 80, 83, 84; ♀ 79; Kilosa, ♂ 81; ♀ 77 mm.

Range.—Central Kenya from Mount Kenya area, Nairobi area, and Ukamba to Kilimanjaro and south through central Tanganyika to Kilosa.

Remarks.—These are the Kenya birds that have usually gone under the name *hamatus* Hartlaub, which, however, applies only to birds from farther south. All the females whose measurements are listed have a black spot in front of the eye, but the spot is reduced in the Kilosa bird, indicating intergradation with more southern forms.

***Dryoscopus cubla hamatus* Hartlaub, 1863**

Like *nairobiensis* except that the female has no black in front of the eye. The range of *hamatus* as here understood is: south and southeastern shores of Lake Victoria south through western and southern Tanganyika Territory to the Zambezi in Portuguese East Africa and west through the southern half of the Belgian Congo and most of Northern Rhodesia to northern and central Angola.

The type locality of *hamatus* is Kaseh, south of Mwanzi on the southern shore of Lake Victoria, and being within the range of birds lacking a black spot in front of the eye (see localities below, under *Measurements*), the name *hamatus* belongs to this type of bird. Another name, *suahelicus* Neumann (1899, Jour. f. Orn., 47: 414), is a synonym. Neumann had specimens from Kakoma and Ugalla (east of Lake Tanganyika and just south of Kaseh), Masai Land, Kilimanjaro and Fort Smith—a mixed series of both *hamatus* and *nairobiensis*. Neumann diagnosed *suahelicus* as differing from Angola birds (*occidentalis*=*hamatus*) on size only (wing 77–81 mm. vs. 84–86 for the latter) and states that the 12 examples from Kakoma and Ugalla have this size; as the other specimens are larger, thus being referred specimens not agreeing precisely with the diagnosis, it seems advisable to restrict the type locality to either Kakoma or Ugalla. I choose Kakoma and consider *suahelicus* a synonym of *hamatus*.

Though Neumann mentioned variations in size in his series, our specimens show little.

Wing measurements.—Kenya: Kericho, ♂ 82–87; ♀ 82. Tanganyika Territory: Mwanza, ♂ 82; ♀ 80. Belgian Congo: Haut Lua-

pula, ♂ 83, 86; ♀ 83, 84. Northern Rhodesia: Lundazi, ♂ 81-87; ♀ 78-85. Angola (northern and central): ♂ 84-90; ♀ 80, 85 mm.

The slight average larger size of Angola birds does not necessitate separating them as *occidentalis* Neumann, 1899.

Dryoscopus cubla okavangensis Roberts, 1932

D. c. okavangensis, type locality Maun, is a quite distinct race, as shown by the gray in the forehead and in the back of our seven Bechuanaland females, compared with more northern races (wing: ♂ 80-87; ♀ 81-82). Three female specimens in the American Museum from Capelongo and Luvando, southern Angola, agree well with Bechuanaland females. Three females from South West Africa (wing 78, 79, 80), though slightly smaller, are also similar.

The range is thus northern Bechuanaland (and probably adjacent Northern Rhodesia) to South West Africa and southern Angola.

Dryoscopus cubla chapini Clancey, 1954

I have not seen this bird from eastern Transvaal. It is apparently like *cubla* but with a red iris.

Dryoscopus cubla cubla Shaw, 1809

D. c. cubla, of South Africa, is said to be separable from all the above on the yellow, not red iris; both *chapini* and *cubla* differ from the other races in having the rump and flanks tinged brownish in the female. One female seen.



UNIVERSITY OF ILLINOIS-URBANA



3 0112 027924254